

Comment on “On the Renormalization of Conductance in Tomonaga-Luttinger Liquid”

Satoshi Fujimoto¹ and Norio Kawakami^{2,3}

¹*Department of Physics, Kyoto University, Kyoto 606*

²*Department of Applied Physics, Osaka University, Suita, Osaka 565*

³*Department of Material and Life Science, Osaka University, Suita, Osaka 565*

Recently Kawabata found that the conductance in Tomonaga-Luttinger (TL) liquids should not be renormalized if one incorporates the renormalization of applied fields correctly. This claim is generalized to include the interactions which carry the large momentum transfer. We thus find that Kawabata’s conclusion holds generally for the class of TL liquids. A key observation is that such irrelevant interactions merely renormalize the velocity and the TL parameter at the massless TL fixed point.

In the recent paper, Kawabata has shown that the conductance in Tomonaga-Luttinger (TL) liquids is not renormalized, i.e. $G = 2e^2/h$, if one correctly takes into account the renormalization of applied fields self-consistently [1]. This is a plausible scenario for the observed conductance [2], which is distinct from the previous proposals [3]. He has demonstrated this fact by using a special version of the TL model. He has also raised a question how his conclusion would be modified if the interactions carrying the large momentum transfer are incorporated, but this problem remains still open. In this comment, we show that Kawabata’s results for the conductance hold even if such interactions are incorporated, and his conclusion is quite general for the TL liquids.

Let us start with the model studied by Kawabata [1],

$$\begin{aligned}\mathcal{H}_0 &= \frac{\hbar^2}{2m} \int dx \psi^\dagger(x) \frac{d^2}{dx^2} \psi(x), \\ \mathcal{H}_1 &= \frac{g}{2\pi} \int dx : \rho(x) \rho(x) :, \end{aligned} \quad (1)$$

where $\psi(x)$ is the Fermi field, $::$ denotes the normal ordering, and $\rho(x) = \rho_+(x) + \rho_-(x)$ with $\rho_+(x)$ ($\rho_-(x)$) being the density of left- (right-) going electrons. This is a special version of the TL model. For simplicity, we have omitted the spin index since only the charge sector is relevant to the conductance. In this translationally invariant model, the renormalization of the Fermi velocity v_F is uniquely related to the TL parameter γ via $v_F \rightarrow v_F/\gamma$. This makes the analysis very simple, and the self-consistent treatment [1] turns out to be exact.

If we take into account more general interactions, the velocity and the TL parameter are renormalized in different manners. We consider the general model describing TL liquids for which the interaction term is given by,

$$\begin{aligned}\mathcal{H}_1 &= \frac{1}{2\pi} \int dx [g_4 \sum_{\alpha=\pm} : \rho_\alpha(x) \rho_\alpha(x) : \\ &\quad + 2g_2 : \rho_+(x) \rho_-(x) :] + \mathcal{H}_U, \end{aligned} \quad (2)$$

where $\mathcal{H}_U$ denotes the interactions other than forward scatterings, which may include the large momentum

transfer. Though it seems difficult at first glance to study the effect of the interactions (2) on the conductance, this problem can be resolved in the following way. The first key observation is that $\mathcal{H}_U$ should become irrelevant after the renormalization so far as the system remains as TL liquids. Hence, what is caused by $\mathcal{H}_U$ is just to renormalize the velocity $v_F \rightarrow v_F^*$ and the interactions $g_2(g_4) \rightarrow g_2^*(g_4^*)$ in different ways. This is a characteristic feature which does not appear in the special model (1) [1]. The renormalized Hamiltonian still has two different types of the interactions g_2^* and g_4^* . For our purpose, it is crucial to rewrite the renormalized Hamiltonian in a suitable form to apply the analysis of Kawabata. Namely, passing to the continuum limit, the renormalized Hamiltonian including the kinetic-energy term is shown to be equivalent to

$$\begin{aligned}\tilde{\mathcal{H}} &= \hbar \tilde{v}_F \sum_{\alpha=\pm} \int dx : \psi_\alpha^\dagger (i\alpha \frac{d}{dx}) \psi_\alpha(x) : \\ &\quad + \frac{\tilde{g}}{2\pi} \int dx : \rho(x) \rho(x) :, \end{aligned} \quad (3)$$

where $\psi_\pm^\dagger$ is the right- (left-) going Fermi fields, $\hbar \tilde{v}_F = \hbar v_F^* + (g_4^* - g_2^*)/\pi$ and $\tilde{g} = g_2^*$. Here we have used the fact that the g_4^* term merely renormalizes the velocity. One can now see that the reduced Hamiltonian (3) has the effective interaction which is translationally invariant, and is essentially the same as (1) studied by Kawabata, so that we can directly follow his arguments on the renormalization of applied fields [4]. Exploiting his results, we thus end up with the conclusion that the conductance in TL liquids is not renormalized, i.e.

$$G = 2e^2/h, \quad (4)$$

even if we take into account the interactions (2) which may carry the large momentum transfer. Therefore Kawabata’s conclusion holds not only for the special model (1) but also for the class of TL liquids generally.

- [1] A. Kawabata: J. Phys. Soc. Jpn. **65** (1996) 30.
- [2] S. Tarucha, T. Honda and T. Saku: Solid State Commun. **94** (1995) 413.
- [3] D. L. Maslov and M. Stone: Phys. Rev. B **52** (1995) R5539; V.V. Ponomarenko: Phys. Rev. B **52** (1995) R8666; H. Safi and H. J. Schulz: Phys. Rev. B **52** (1995) R17040.
- [4] The TL parameter is given by $\gamma = [1 + \tilde{g}/(\pi\hbar\tilde{v}_F)]^{-1/2}$.